


Question 1

(9 marks)

- (a) A sailing boat leaves port and sails on a bearing of 300° for 4 km before turning to sail 1.9 km on a bearing of 080° . How far is the boat from the port and what is the bearing of final position of the boat from the port. (3 marks)



$$r = \sqrt{4^2 + 1.9^2 - 2 \times 4 \times 1.9 \cos 40^\circ}$$

$$= 2.82 \text{ km}$$

$$\frac{\sin \theta}{1.9} = \frac{\sin 40^\circ}{2.82}$$

$$\theta = \sin^{-1} \left(\frac{1.9 \times \sin 40^\circ}{2.82} \right) = 25.64^\circ$$


Bearing of the boat (B) from the port (P) is $300^\circ + 25.64^\circ$
 $\approx 326^\circ \text{ T}$

✓ Uses cosine rule

✓ Uses sine rule

✓ Correct bearing

- (b) Find the magnitude and direction of the resultant of the pair of forces in the diagram below. (3 marks)



$$|\vec{r}| = \sqrt{7^2 + 3^2 - 2 \times 7 \times 3 \cos 70^\circ}$$

$$= 6.6 \text{ N}$$

$$\frac{\sin \theta}{3} = \frac{\sin 70^\circ}{6.6}$$

$$\theta = \sin^{-1} \left(\frac{3 \times \sin 70^\circ}{6.6} \right) = 25.3^\circ$$


$$\therefore \text{Bearing} = 100^\circ + 25.3^\circ \approx 125^\circ \text{ T}$$

✓ Correct magnitude of resultant vector

✓ Uses sine rule to find θ

✓ Correct bearing

- (c) In still air, an aircraft can maintain a speed of 350 km/h. In what direction should the aircraft be pointing if it wished to travel in a direction 160° , and a 35 km/h wind is blowing from 075° ? (3 marks)



$\vec{a}$ is the velocity of the aircraft.

$\vec{w}$ is the velocity of the wind.

$\vec{r}$ is the resultant velocity.

$$\frac{\sin \theta}{35} = \frac{\sin 95^\circ}{350}$$

$$\theta = \sin^{-1} \left(\frac{35 \times \sin 95^\circ}{350} \right) = 5.7^\circ$$

$$\therefore \text{Bearing of } \vec{a} = 160^\circ - 5.7^\circ \approx 154^\circ \text{ T}$$

✓ Correct diagram
 ✓ Uses sine rule to find θ

✓ Correct bearing

Question 2**(6 marks)**

(a) Vectors **a** and **b** have the same magnitude and vectors **a** and **c** are perpendicular, where

$\mathbf{a} = \begin{bmatrix} m \\ n \end{bmatrix}$, $\mathbf{b} = \begin{bmatrix} -4 \\ 6 \end{bmatrix}$ and $\mathbf{c} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$. Determine the values of m and n . (3 marks)

$$\begin{cases} m^2 + n^2 = (-4)^2 + 6^2 = 52 \\ 2m + 3n = 0 \end{cases}$$


$$\therefore \begin{cases} m = -6 \\ n = 4 \end{cases} \quad \begin{cases} m = 6 \\ n = -4 \end{cases}$$

✓ Correct equations of m and n

✓ Correct values of m

✓ Correct values of n

(b) Determine the scalar projection of a velocity of 12 m/s on a bearing of 065° onto a velocity of 20 m/s on a bearing of 280° , giving your answer to two decimal places.

(3 marks)

$$\begin{aligned} & -12 \times \cos(180^\circ - 80^\circ - 65^\circ) \quad \text{or} \quad 12 \times \cos(80^\circ + 65^\circ) \\ & = -12 \times \cos 35^\circ \quad \text{or} \quad 12 \times \cos 145^\circ \\ & = -9.83 \end{aligned}$$

✓ correct angle

✓ correct direction (negative)

✓ correct scalar projection

Question 3

(8 marks)

(a) A triangle has vertices as $A(-3, 1)$, $B(-1, 4)$ and $C(5, 0)$.

(i) Determine the vectors $\vec{AB}$, $\vec{AC}$ and $\vec{BC}$.

(2 marks)

$$\vec{AB} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$$

$$\vec{AC} = \begin{bmatrix} 8 \\ -1 \end{bmatrix}$$

$$\vec{BC} = \begin{bmatrix} 6 \\ -4 \end{bmatrix}$$

-1 each error

(ii) Use a vector method to prove that triangle ABC is right-angle.

(2 marks)

$$\vec{AB} \cdot \vec{BC} = \begin{bmatrix} 2 \\ 3 \end{bmatrix} \cdot \begin{bmatrix} 6 \\ -4 \end{bmatrix}$$

$$= 12 - 12$$

$$= 0$$


✓ Expresses
 $\vec{AB} \cdot \vec{BC}$

✓ Evaluates
 $\vec{AB} \cdot \vec{BC}$

$\therefore \vec{AB} \perp \vec{BC} \Rightarrow \triangle ABC$ right-angled at B.

(b) Use a vector method to prove that if the diagonals of a parallelogram are perpendicular to each other, then the parallelogram is a rhombus.

(4 marks)



$$\text{Let } \vec{OA} = \vec{a}, \vec{OB} = \vec{b},$$

$$\text{so } \vec{OC} = \vec{a} + \vec{b}, \vec{AB} = \vec{b} - \vec{a}$$

If $\vec{OC}$ & $\vec{AB}$ are perpendicular, then

$$\vec{OC} \cdot \vec{AB} = 0$$

$$(\vec{a} + \vec{b}) \cdot (\vec{b} - \vec{a}) = 0$$

$$\vec{a} \cdot \vec{b} - \vec{a} \cdot \vec{a} + \vec{b} \cdot \vec{b} - \vec{a} \cdot \vec{b} = 0$$

✓ States $\vec{OC} \cdot \vec{AB} = 0$

✓ States $\vec{OA} \cdot \vec{OA} = \vec{OB} \cdot \vec{OB}$

✓ States $|\vec{OA}|^2 = |\vec{OB}|^2$

✓ States $|\vec{OA}| = |\vec{OB}|$

$$\vec{a} \cdot \vec{a} = \vec{b} \cdot \vec{b}$$

$$\therefore |\vec{a}|^2 = |\vec{b}|^2$$

$$\therefore |\vec{a}| = |\vec{b}|$$


$$\therefore |\vec{OA}| = |\vec{OB}|$$

Hence, $\square OACB$ is a rhombus

Question 4


(8 marks)

Two forces, $F_1 = 550\text{N}$ and $F_2 = 770\text{N}$, act on a body at O, and make angles of $\alpha = 33^\circ$, and $\beta = 18^\circ$ respectively with the vertical OP, as shown in the diagram below.



(a) Determine the magnitude of resultant force and the angle it makes with the vertical.

(5 marks)



$$F_R = \sqrt{550^2 + 770^2 - 2 \times 550 \times 770 \times \cos(180 - 33 - 18)}$$

$$= 1195.17\text{N}$$

$$\frac{\sin \theta}{770} = \frac{\sin 129^\circ}{1195.17}$$

$$\theta = 30^\circ$$

$$\text{Required Angle} = 33^\circ - 30^\circ = 3^\circ$$

✓ Sketch with force nose to tail

✓ Indicate use of cosine rule for magnitude

✓ Correct magnitude


✓ Indicate use of sine rule for angle

✓ Angle with vertical

(b) The magnitude of F_1 is to be adjusted so that the direction of the resultant is vertical.

Determine the required magnitude of F_1 .

(3 marks)



$$\frac{\sin \alpha}{F_2} = \frac{\sin \beta}{F_1}$$

$$F_1 = \frac{770 \times \sin 18^\circ}{\sin 33^\circ} = 436.88\text{N (2 d.p.)} \approx 437\text{N}$$

✓ Sketch

✓ Indicate use of sine rule


✓ Correct magnitude

Question 5

(9 marks)

(a) The work done, in joules, by a force of $\mathbf{F}$ Newtons in changing the displacement of an object by $\mathbf{s}$ metres, is given by the scalar product of $\mathbf{F}$ and $\mathbf{s}$.

(i) A force of 250 N acting due south moves an object 4.3 m in a south-westerly direction. Determine the work done. (2 marks)



$$\begin{aligned} W &= \vec{s} \cdot \vec{F} \\ &= |\vec{s}| |\vec{F}| \cos 45^\circ \\ &= 4.3 \times 250 \times \cos 45^\circ \\ &= 760 \text{ J} \end{aligned}$$

✓ Substitute correctly
✓ Evaluate correctly


(ii) Another force of 155 N does 269 joules of work in moving an object 190 cm. Determine the angle between the force and the direction of movement. (2 marks)

$$\begin{aligned} 155 \times 1.9 \times \cos \theta &= 269 \\ \theta &= \cos^{-1} \left(\frac{269}{155 \times 1.9} \right) \\ &= 24^\circ \end{aligned}$$

✓ Substitute correctly
✓ Evaluate correctly

(b) A triangle is formed by three non-zero vectors $\mathbf{a}$, $\mathbf{b}$ and $\mathbf{c}$, so that $\mathbf{c} = \mathbf{a} - \mathbf{b}$, and θ is the angle between $\mathbf{a}$ and $\mathbf{b}$.

(i) Sketch the triangle. (1 mark)



✓ Sketch

(ii) Explain why $\mathbf{c} \cdot \mathbf{c} = |\mathbf{c}|^2$. (1 mark)

$$\begin{aligned} \vec{c} \cdot \vec{c} &= |\vec{c}| |\vec{c}| \cos \theta \\ \text{Since } \theta &= 0^\circ, \cos \theta = 1 \\ \text{Hence, } \vec{c} \cdot \vec{c} &= |\vec{c}|^2 \end{aligned}$$

✓ Explanation

(iii) Use $\mathbf{c} \cdot \mathbf{c} = (\mathbf{a} - \mathbf{b}) \cdot (\mathbf{a} - \mathbf{b})$ to deduce the cosine rule. (3 marks)

$$\begin{aligned} \vec{c} \cdot \vec{c} &= (\vec{a} - \vec{b}) \cdot (\vec{a} - \vec{b}) \\ \vec{c} \cdot \vec{c} &= \vec{a} \cdot \vec{a} - \vec{a} \cdot \vec{b} - \vec{b} \cdot \vec{a} + \vec{b} \cdot \vec{b} \\ \therefore |\vec{c}|^2 &= |\vec{a}|^2 - |\vec{a}| |\vec{b}| \cos \theta - |\vec{b}| |\vec{a}| \cos \theta + |\vec{b}|^2 \\ \text{Hence, } |\vec{c}|^2 &= |\vec{a}|^2 + |\vec{b}|^2 - 2|\vec{a}| |\vec{b}| \cos \theta \end{aligned}$$

✓ Expand scalar product
✓ Use $\vec{c} \cdot \vec{c} = |\vec{c}|^2$ from (ii)
✓ Use scalar product definition

Question 6


(11 marks)

A small boat that can maintain a steady speed of 5 ms^{-1} is to cross a river from A to B , where $\vec{AB} = (35\mathbf{i} - 105\mathbf{j}) \text{ m}$.

A current of $(-\mathbf{i} - 2\mathbf{j}) \text{ ms}^{-1}$ flows in the river.

The velocity vector that the pilot of the small boat must set to travel from A to B is $a\mathbf{i} + b\mathbf{j}$, where a and b are constants.

(a) Explain why $t(a - 1) = 35$ and $t(b - 2) = -105$, where t is a constant. (3 marks)



Resultant velocity $\vec{v}_R = \vec{v} + \vec{c} = (a\mathbf{i} + b\mathbf{j}) + (-\mathbf{i} - 2\mathbf{j})$
 $= (a-1)\mathbf{i} + (b-2)\mathbf{j}$ is parallel to $\vec{AB}$

$\therefore \vec{AB} = \vec{v}_R t = ((a-1)\mathbf{i} + (b-2)\mathbf{j})t$

$= t(a-1)\mathbf{i} + t(b-2)\mathbf{j} = 35\mathbf{i} - 105\mathbf{j}$

Hence, $t(a-1) = 35$ & $t(b-2) = -105$

(b) Eliminate t from the equations in (a) and hence express b in terms of a , simplifying your expression. (3 marks)

$\therefore t(a-1) = 35 \Rightarrow t = \frac{35}{a-1}$ & $t(b-2) = -105 \Rightarrow t = \frac{-105}{b-2}$

$\therefore \frac{35}{a-1} = \frac{-105}{b-2}$

$35b - 70 = -105a + 105$

$b - 2 = -3a + 3$

$b = -3a + 5$

✓ Eliminate t

✓ Cross multiply

✓ Simplify

(c) Explain why $a^2 + b^2 = 25$.

(1 mark)

The magnitude of $\vec{v} = a\mathbf{i} + b\mathbf{j}$ is $\sqrt{a^2 + b^2} = \sqrt{25}$

Hence the speed of the small boat is 5 m/s

✓ State magnitude and speed

- (d) Use your equations from (b) and (c) to determine the values of a and b . (3 marks)

$$\begin{cases} b = -3a + 5 \\ a^2 + b^2 = 25 \end{cases} \Rightarrow a^2 + (-3a + 5)^2 = 25$$

$$\Rightarrow a = 3 \text{ or } a = 0 (\text{exclude})$$

$$\therefore b = -3 \times 3 + 5 = -4$$

$$\therefore \vec{v} = 3\vec{i} - 4\vec{j}$$

✓ Correct equation

✓ Solve a & b

✓ Eliminate alternative solution

- (e) Determine the time that the small boat will take to travel from A to B . (1 mark)

$$t = \frac{35}{a-1} = \frac{35}{2} = 17.5s$$

✓ Correct time

Question 7

(8 marks)

Three vectors are given by $\mathbf{a} = 3\mathbf{i} - 4\mathbf{j}$, $\mathbf{b} = -3\mathbf{i} + 1.5\mathbf{j}$ and $\mathbf{c} = -2\mathbf{i} + y\mathbf{j}$, where y is a constant.

- (a) Determine the vector projection of $\mathbf{b}$ on $\mathbf{a}$.

(3 marks)

$$\vec{b} \cdot \vec{a} = 3 \times (-3) + (-4) \times (1.5) = -15$$

$$\vec{a} \cdot \vec{a} = 3^2 + (-4)^2 = 25$$

$$\begin{aligned} \vec{u} &= \frac{\vec{b} \cdot \vec{a}}{\vec{a} \cdot \vec{a}} \vec{a} = \frac{-15}{25} \times (3\vec{i} - 4\vec{j}) \\ &= -\frac{9}{5}\vec{i} + \frac{12}{5}\vec{j} \end{aligned}$$

✓ Evaluate $\vec{b} \cdot \vec{a}$ &
 $\vec{a} \cdot \vec{a}$

✓ State $\vec{u} = \frac{\vec{b} \cdot \vec{a}}{\vec{a} \cdot \vec{a}} \vec{a}$

✓ Correct vector projection

- (b) Determine the value(s) of y if

- (i) $\mathbf{a}$ and $\mathbf{c}$ are perpendicular.

(2 marks)

$$\vec{a} \cdot \vec{c} = 0$$

$$3 \times (-2) - 4y = 0$$

$$y = -\frac{3}{2}$$

✓ Use scalar product

✓ Solve y

- (ii) the angle between the directions of $\mathbf{b}$ and $\mathbf{c}$ is 45° .

(3 marks)

$$\vec{b} \cdot \vec{c} = 6 + 1.5y$$

$$|\vec{b}| = \sqrt{(-3)^2 + 1.5^2} = \frac{3\sqrt{5}}{2}$$

$$|\vec{c}| = \sqrt{(-2)^2 + y^2} = \sqrt{4 + y^2}$$

$$\therefore \vec{b} \cdot \vec{c} = |\vec{b}| |\vec{c}| \cos 45^\circ$$



$$6 + 1.5y = \frac{3\sqrt{5}}{2} \times \sqrt{4 + y^2} \times \frac{\sqrt{2}}{2}$$

$$y = -\frac{2}{3}, y = 6$$

✓ Use scalar product

✓ State one solution

✓ State second solution



$[-3, 1.5] \Rightarrow b$

$[-3 \ 1.5]$

$[-2, y] \Rightarrow c$

$[-2 \ y]$

$\text{norm}(b) \times \text{norm}(c) \times \cos(45) = \text{dotP}(b, c)$

$$0.75 \cdot (10 \cdot (y^2 + 4))^{0.5} = 1.5 \cdot y + 6$$

$\text{solve}(\text{ans}, y, 0, -\infty, \infty)$

$\{y = -0.6666666667, y = 6\}$