

Rossmoyne Senior High School



Semester One Examination, 2018

Question/Answer booklet

MATHEMATICS SPECIALIST UNIT 1

Section One:
Calculator-free

SOLUTIONS

Student Name: _____

Teacher's Name: _____

Time allowed for this section

Reading time before commencing work: five minutes
Working time: fifty minutes

Materials required/recommended for this section

To be provided by the supervisor

This Question/Answer booklet
Formula sheet

To be provided by the candidate

Standard items: pens (blue/black preferred), pencils (including coloured), sharpener,
correction fluid/tape, eraser, ruler, highlighters

Special items: nil

Important note to candidates

No other items may be taken into the examination room. It is **your** responsibility to ensure that you do not have any unauthorised material. If you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

Structure of this paper

Section	Number of questions available	Number of questions to be answered	Working time (minutes)	Marks available	Percentage of examination
Section One: Calculator-free	8	8	50	54	36
Section Two: Calculator-assumed	13	13	100	97	64
Total					100

Instructions to candidates

1. The rules for the conduct of examinations are detailed in the school handbook. Sitting this examination implies that you agree to abide by these rules.
2. Write your answers in this Question/Answer booklet.
3. You must be careful to confine your response to the specific question asked and to follow any instructions that are specified to a particular question.
4. Supplementary pages for the use of planning/continuing your answer to a question have been provided at the end of this Question/Answer booklet. If you use these pages to continue an answer, indicate at the original answer where the answer is continued, i.e. give the page number.
5. Show all your working clearly. Your working should be in sufficient detail to allow your answers to be checked readily and for marks to be awarded for reasoning. Incorrect answers given without supporting reasoning cannot be allocated any marks. For any question or part question worth more than two marks, valid working or justification is required to receive full marks. If you repeat any question, ensure that you cancel the answer you do not wish to have marked.
6. It is recommended that you do not use pencil, except in diagrams.
7. The Formula sheet is not to be handed in with your Question/Answer booklet.

Section One: Calculator-free

36% (54 Marks)

This section has **eight (8)** questions. Answer **all** questions. Write your answers in the spaces provided.

Working time: 50 minutes.

Question 1

(5 marks)

Relative to the origin O , points A and B have position vectors $-2\mathbf{i} - 3\mathbf{j}$ and $\mathbf{i} - 6\mathbf{j}$ respectively.

- (a) Determine the unit vector $\hat{\mathbf{c}}$, where $\mathbf{c} = \overrightarrow{AB}$. (3 marks)

Solution
$\mathbf{c} = \begin{pmatrix} 1 \\ -6 \end{pmatrix} - \begin{pmatrix} -2 \\ -3 \end{pmatrix} = \begin{pmatrix} 3 \\ -3 \end{pmatrix}$
$ \mathbf{c} = \sqrt{9 + 9} = 3\sqrt{2}$
$\hat{\mathbf{c}} = \frac{1}{3\sqrt{2}} \begin{pmatrix} 3 \\ -3 \end{pmatrix} = \frac{\sqrt{2}}{2} \begin{pmatrix} 1 \\ -1 \end{pmatrix}$
Specific behaviours
✓ vector $\mathbf{c}$ ✓ magnitude ✓ unit vector, simplified

- (b) Vector $\mathbf{d}$ has magnitude $4\sqrt{2}$, is parallel to $\mathbf{c}$ and in the opposite direction. Determine $\mathbf{d}$. (2 marks)

Solution
$\mathbf{d} = 4\sqrt{2} \times (-1) \times \frac{\sqrt{2}}{2} \begin{pmatrix} 1 \\ -1 \end{pmatrix}$ $= \begin{pmatrix} -4 \\ 4 \end{pmatrix}$
Specific behaviours
✓ reverses $\mathbf{c}$ ✓ correct vector

Question 2

(5 marks)

Let the displacement vectors **a**, **b** and **c** be $(11, -4)$, $(5, 14)$ and $(8, m)$ respectively, where m is a real number.

- (a) Determine the vector $3\mathbf{a} + 2\mathbf{b}$.

(2 marks)

Solution
$3\mathbf{a} + 2\mathbf{b} = 3(11, -4) + 2(5, 14)$ $= (33, -12) + (10, 28)$ $= (43, 16)$
Specific behaviours
✓ multiplies by scalar ✓ correct vector

- (b) Given that $|\mathbf{a} + \mathbf{b} + k\mathbf{c}| = 0$, determine the values of k and m .

(3 marks)

Solution
$\begin{pmatrix} 11 \\ -4 \end{pmatrix} + \begin{pmatrix} 5 \\ 14 \end{pmatrix} + k \begin{pmatrix} 8 \\ m \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$ From i-coeff: $11 + 5 + 8k = 0 \Rightarrow k = -2$ From j-coeff: $-4 + 14 - 2m = 0 \Rightarrow m = 5$
Specific behaviours
✓ vector equation ✓ value of k ✓ value of m

Question 3

(8 marks)

Consider the following statement:

If a polygon has an interior angle sum of 360° , then it is a square.

- (a) Use a **counter-example** to explain why the statement is false.

(2 marks)

Solution
A trapezium has an interior angle sum of 360° but is not a square.
Specific behaviours
✓ names or draws any quadrilateral that is not a square ✓ uses angle sum and fact that shape is not a square

- (b) Write the **converse** statement and state whether it is always, sometimes or never true.

(2 marks)

Solution
If it is a square, then it has an interior angle sum of 360°. The converse statement is always true.
Specific behaviours
✓ writes converse ✓ states always true

- (c) Write the **inverse** statement and state whether it is always, sometimes or never true.

(2 marks)

Solution
If it does not have an interior angle sum of 360°, then it is not a square. The inverse statement is always true.
Specific behaviours
✓ writes inverse ✓ states always true

- (d) Write the **contrapositive** statement and state whether it is always, sometimes or never true.

(2 marks)

Solution
If it is not a square, then it does not have an interior angle sum of 360°. The contrapositive statement is sometimes true. (eg true for triangle, false for any quadrilateral)
Specific behaviours
✓ writes contrapositive ✓ states sometimes true

Question 4

(6 marks)

- (a) Determine the value of the constant n , given that the vectors $5\mathbf{i} + 3\mathbf{j}$ and $-8\mathbf{i} + n\mathbf{j}$ are perpendicular. (2 marks)

Solution
$-40 + 3n = 0$ $n = \frac{40}{3} = 13\frac{1}{3}$
Specific behaviours
✓ equates scalar product to 0 ✓ solves for n

- (b) The vectors $\mathbf{a}$ and $\mathbf{b}$ are such that $|\mathbf{a}| = 17$, $|\mathbf{b}| = 13$ and $\mathbf{a} \cdot \mathbf{b} = -21$. Evaluate

- (i) $-2\mathbf{a} \cdot 5\mathbf{b}$. (1 mark)

Solution
$-21 \times -2 \times 5 = 210$
Specific behaviours
✓ correct value


- (ii) $(\mathbf{b} + \mathbf{a}) \cdot (\mathbf{a} - \mathbf{b})$. (3 marks)

Solution
$(\mathbf{b} + \mathbf{a}) \cdot (\mathbf{a} - \mathbf{b}) = \mathbf{a} \cdot \mathbf{a} - \mathbf{b} \cdot \mathbf{b}$ $= \mathbf{a} ^2 - \mathbf{b} ^2$ $= 17^2 - 13^2$ $= (17 + 13)(17 - 13)$ $= 30 \times 4 = 120$
Specific behaviours
✓ expands ✓ recognises $ \mathbf{a} ^2 = a^2$ and $ \mathbf{b} ^2 = b^2$ ✓ correct value

Question 5

(8 marks)

- (a) In the diagram below, the vertices of triangle ABC lie on a circle with centre O . Given that $\angle ABC = 54^\circ$, determine the values of $\angle AOC$ and $\angle OAC$. (2 marks)



Solution
$\angle AOC = 2 \times 54 = 108^\circ$
$\angle OAC = \frac{180 - 108}{2} = 36^\circ$
Specific behaviours
✓ first angle
✓ second angle


- (b) Prove that the size of the angle subtended by an arc at the centre of a circle is twice the size of the angle subtended at any point on the circumference by the same arc. Assume only basic axioms and properties of triangles can be used. (6 marks)

Solution
Required to prove that $a = 2b$ Let $b = \angle OQP + \angle OQR$ But $\angle OQP = \angle OPQ$ and $\angle OQR = \angle ORQ$ (isosceles triangles) And so $\angle QOP = 180 - 2\angle OQP$ and $\angle QOR = 180 - 2\angle OQR$ At O, $a = 360 - \angle QOP - \angle QOR$ $a = 360 - (180 - 2\angle OQP) - (180 - 2\angle OQR)$ $a = 2(\angle OQP + \angle OQR)$ $a = 2b$, as required.
Specific behaviours
✓ labelled diagram(s) illustrating RTP ✓✓ congruence statement with appropriate reason ✓ expressions for angles at O ✓ equation using angle sum at a point ✓ substitutes and simplifies


Question 6**(6 marks)**

A drone leaves point P and travels 74 m on bearing of 290° to Q , then 40 m on bearing 020° to R and finally 114 m on bearing 110° to S .

- (a) Sketch a neat diagram to show the path of the drone.

(2 marks)


- (b) The drone is to return directly from S to P . Determine the distance it must fly and on what bearing. **(4 marks)**



Question 7

(7 marks)

A parallelogram OABC has D the midpoint of AB, E the midpoint of BC and DC intersecting OE at F. Let $\vec{OA} = \mathbf{a}$ and $\vec{OC} = \mathbf{c}$.



a) Determine the following in terms of $\mathbf{a}$ and $\mathbf{c}$:

(2 marks)

i. $\vec{OE}$

Solution
$\vec{OE} = \vec{OC} + \frac{1}{2}\vec{CB} = \mathbf{c} + \frac{1}{2}\mathbf{a}$
Specific behaviours
✓ correctly state $\vec{OE}$ in terms of $\mathbf{a}$ and $\mathbf{c}$

ii. $\vec{DC}$

Solution
$\vec{DC} = \vec{DO} + \vec{OC} = -\frac{1}{2}\mathbf{c} - \mathbf{a} + \mathbf{c} = -\mathbf{a} + \frac{1}{2}\mathbf{c}$
Specific behaviours
✓ correctly state $\vec{DC}$ in terms of $\mathbf{a}$ and $\mathbf{c}$

b) By letting $\vec{DF} = h\vec{DC}$ and $\vec{OF} = k\vec{OE}$, show that $h = \frac{3}{5}$ and $k = \frac{4}{5}$.

(5 marks)

Solution
$\vec{DF} = \frac{h}{2}\mathbf{c} - h\mathbf{a}$ $\vec{OF} = k\mathbf{c} + \frac{k}{2}\mathbf{a}$
$\vec{OF} = \vec{OD} + \vec{DF} \Rightarrow k\mathbf{c} + \frac{k}{2}\mathbf{a} = \frac{1}{2}\mathbf{c} + \mathbf{a} + \frac{h}{2}\mathbf{c} - h\mathbf{a}$
$(k - \frac{1}{2} - \frac{h}{2})\mathbf{c} = (1 - h - \frac{k}{2})\mathbf{a}$
As $\mathbf{a}$ not P $\mathbf{c}$, $k - \frac{1}{2} - \frac{h}{2} = 0$ and $1 - h - \frac{k}{2} = 0 \Rightarrow h = \frac{3}{5}$ and $k = \frac{4}{5}$
Specific behaviours
✓ Expresses $\vec{DF}$ and $\vec{OF}$ in terms of $\mathbf{a}$ and $\mathbf{c}$ ✓ States and equation linking $\vec{OF}$, $\vec{DF}$ and $\vec{OD}$ ✓ Rearranges equation into $h\mathbf{a} = k\mathbf{c}$ form ✓ Uses $h = 0$ and $k = 0$ ✓ Correctly solves for h and k

Question 8

(9 marks)

- (a) Evaluate
- ${}^{25}P_{19} \div {}^{23}P_{20}$
- .

(3 marks)

Solution
$\frac{25!}{6!} \times \frac{3!}{23!} = \frac{25 \times 24}{6 \times 5 \times 4} = 5$
Specific behaviours
✓ expresses using factorials ✓ eliminates factorials ✓ evaluates

- (b) Express
- $8! + 7! + 6!$
- in the form
- $a^2b!$
- , where
- a
- and
- b
- are positive integers.

(3 marks)

Solution
$8! + 7! + 6! = (8 \times 7 + 7 + 1) \times 6!$ $= 64 \times 6!$ $= 8^2 \times 6!$
Specific behaviours
✓ factors out lowest factorial ✓ simplifies ✓ writes in required form

- (c) Show that for
- $n \in \mathbb{Z}, n \geq 1$
- , the sum
- $(n+2)! + (n+1)! + n!$
- can always be expressed in the form
- $a^2b!$
- where
- a
- and
- b
- are positive integers.

(3 marks)

Solution
$(n+2)! + (n+1)! + n! = ((n+2)(n+1) + (n+1) + 1)n!$ $= (n^2 + 4n + 4)n!$ $= (n+2)^2n!$ $(a = n+2 \text{ and } b = n)$
Specific behaviours
✓ uses $n!$ as one factor ✓ clearly shows composition of second factor ✓ simplifies second factor and writes as required

Supplementary page

Question number: _____

