



Rossmoyne Senior High School

Semester One Examination, 2020

Question/Answer booklet

MATHEMATICS SPECIALIST UNIT 1

Section One:
Calculator-free

SOLUTIONS

WA student number: In figures

--	--	--	--	--	--	--	--

In words

Your name

Time allowed for this section

Reading time before commencing work: five minutes
Working time: fifty minutes

Number of additional
answer booklets used
(if applicable):

--

Materials required/recommended for this section

To be provided by the supervisor

This Question/Answer booklet
Formula sheet

To be provided by the candidate

Standard items: pens (blue/black preferred), pencils (including coloured), sharpener,
correction fluid/tape, eraser, ruler, highlighters

Special items: nil

Important note to candidates

No other items may be taken into the examination room. It is **your** responsibility to ensure that you do not have any unauthorised material. If you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

Structure of this paper

Section	Number of questions available	Number of questions to be answered	Working time (minutes)	Marks available	Percentage of examination
Section One: Calculator-free	8	8	50	52	35
Section Two: Calculator-assumed	13	13	100	98	65
Total					100

Instructions to candidates

1. The rules for the conduct of examinations are detailed in the school handbook. Sitting this examination implies that you agree to abide by these rules.
2. Write your answers in this Question/Answer booklet preferably using a blue/black pen. Do not use erasable or gel pens.
3. You must be careful to confine your answers to the specific question asked and to follow any instructions that are specific to a particular question.
4. Show all your working clearly. Your working should be in sufficient detail to allow your answers to be checked readily and for marks to be awarded for reasoning. Incorrect answers given without supporting reasoning cannot be allocated any marks. For any question or part question worth more than two marks, valid working or justification is required to receive full marks. If you repeat any question, ensure that you cancel the answer you do not wish to have marked.
5. It is recommended that you do not use pencil, except in diagrams.
6. Supplementary pages for planning/continuing your answers to questions are provided at the end of this Question/Answer booklet. If you use these pages to continue an answer, indicate at the original answer where the answer is continued, i.e. give the page number.
7. The Formula sheet is not to be handed in with your Question/Answer booklet.

Section One: Calculator-free

35% (52 Marks)

This section has **eight** questions. Answer **all** questions. Write your answers in the spaces provided.

Working time: 50 minutes.

Question 1

(5 marks)

- (a) Without prefacing the statement with 'It is not ...' or similar, write the negation of the following statements:

- (i) Polygon X is a pentagon or a hexagon.

(1 mark)

Solution
Polygon X is not a pentagon and not a hexagon.
Specific behaviours
✓ correct negation

- (ii) $\forall$ obtuse angles α , the angle 2α is a reflex angle.

(2 marks)

Solution
$\exists$ an obtuse angle α such that 2α is not a reflex angle.
Specific behaviours
✓ replaces 'for all' with 'there exists'
✓ correct negation

- (b) If a true statement is negated, explain whether the contrapositive of the negated statement will be false.

(2 marks)

Solution
The contrapositive of the negated statement will be false. When the true statement is negated, this results in a statement which is false. When the contrapositive of this is taken, the resulting statement will remain false.
Specific behaviours
✓ states that the contrapositive of the negated statement is false, explaining that the negated statement is false
✓ explains that the contrapositive statement is false

Question 2

(6 marks)

The position vectors of points A and B are $\mathbf{r}_A = \begin{pmatrix} -4 \\ 11 \end{pmatrix}$ and $\mathbf{r}_B = \begin{pmatrix} 2 \\ -13 \end{pmatrix}$.

- (a) Determine the position vector of point P that divides AB internally in the ratio 1:5.

(3 marks)

Solution
$\overrightarrow{AB} = \begin{pmatrix} 2 \\ -13 \end{pmatrix} - \begin{pmatrix} -4 \\ 11 \end{pmatrix} = \begin{pmatrix} 6 \\ -24 \end{pmatrix}$ $P = A + \frac{1}{6}\overrightarrow{AB}$ $= \begin{pmatrix} -4 \\ 11 \end{pmatrix} + \frac{1}{6}\begin{pmatrix} 6 \\ -24 \end{pmatrix}$ $= \begin{pmatrix} -3 \\ 7 \end{pmatrix}$
Specific behaviours
✓ vector $\overrightarrow{AB}$ ✓ indicates appropriate method ✓ correct position vector

- (b) A small body leaves B and moves with a constant velocity in a direction parallel to $\begin{pmatrix} -3 \\ 2 \end{pmatrix}$. Determine, with reasons, whether the body will pass through point C with position vector $\mathbf{r}_C = \begin{pmatrix} -20 \\ 1 \end{pmatrix}$.

(3 marks)

Solution
$\overrightarrow{BC} = \begin{pmatrix} -20 \\ 1 \end{pmatrix} - \begin{pmatrix} 2 \\ -13 \end{pmatrix} = \begin{pmatrix} -22 \\ 14 \end{pmatrix}$ Note that $\begin{pmatrix} -22 \\ 14 \end{pmatrix} \neq k \begin{pmatrix} -3 \\ 2 \end{pmatrix}$ and so body will not pass through C as $\overrightarrow{BC}$ is not parallel to $\begin{pmatrix} -3 \\ 2 \end{pmatrix}$.
Specific behaviours
✓ vector $\overrightarrow{BC}$ ✓ indicates $\overrightarrow{BC}$ not parallel to velocity ✓ states will not pass through C

Question 3

(7 marks)

Two forces are given by $\mathbf{F}_1 = 5\mathbf{i} - 2\mathbf{j}$ N and $\mathbf{F}_2 = -\mathbf{i} + 3\mathbf{j}$ N.

(a) Determine

(i) $\mathbf{F}_1 - \mathbf{F}_2$.

(1 mark)

Solution
$\begin{pmatrix} 5 \\ -2 \end{pmatrix} - \begin{pmatrix} -1 \\ 3 \end{pmatrix} = \begin{pmatrix} 6 \\ -5 \end{pmatrix}$ N
Specific behaviours
✓ correct vector

(ii) $2\mathbf{F}_1 + 4\mathbf{F}_2$.

(2 marks)

Solution
$2\begin{pmatrix} 5 \\ -2 \end{pmatrix} + 4\begin{pmatrix} -1 \\ 3 \end{pmatrix} = \begin{pmatrix} 10 \\ -4 \end{pmatrix} + \begin{pmatrix} -4 \\ 12 \end{pmatrix} = \begin{pmatrix} 6 \\ 8 \end{pmatrix}$ N
Specific behaviours
✓ one correct multiple
✓ correct vector

(iii) $|\mathbf{F}_1|$.

(1 mark)

Solution
$\left \begin{pmatrix} 5 \\ -2 \end{pmatrix} \right = \sqrt{(5)^2 + (-2)^2} = \sqrt{29}$ N
Specific behaviours
✓ correct value

(b) The resultant of $5\mathbf{F}_1$, $10\mathbf{F}_2$ and a third force is $12\mathbf{i} + 25\mathbf{j}$ N. Determine the magnitude of the third force. (3 marks)

Solution
$\mathbf{F}_3 = \begin{pmatrix} 12 \\ 25 \end{pmatrix} - \begin{pmatrix} 25 \\ -10 \end{pmatrix} - \begin{pmatrix} -10 \\ 30 \end{pmatrix}$ $= \begin{pmatrix} -3 \\ 5 \end{pmatrix}$
$ \mathbf{F}_3 = \sqrt{9 + 25} = \sqrt{34}$
Specific behaviours
✓ correct vector equation
✓ correct third force
✓ correct magnitude

Question 4

(6 marks)

(a) Consider the statement $a = 3 \Rightarrow 4a = 12$.

(i) Write the converse statement.

(1 mark)

Solution
$4a = 12 \Rightarrow a = 3$
Specific behaviours
✓ correct converse

(ii) Write the inverse statement.

(1 mark)

Solution
$a \neq 3 \Rightarrow 4a \neq 12$
Specific behaviours
✓ correct inverse

(b) State whether each of the following statements are true or false, supporting each answer with an example or counterexample.

(i) $\forall x \in \{\mathbb{R}: x \geq 0\}, \sqrt{x} \leq x$.

(2 marks)

Solution
False. If $x = 0.25$ then $\sqrt{x} = \sqrt{0.25} = 0.5$ but $0.25 < 0.5$.
Specific behaviours
✓ states false, with counterexample ✓ counterexample using $0 < x < 1$

(ii) $\exists n \in \mathbb{Z}$ such that $\forall m \in \mathbb{Z}, n + m = 10$.

(2 marks)

Solution
True. If $m = 8$, we can choose $n = 2$ so that $n + m = 10$.
Specific behaviours
✓ states true, with example ✓ example using integers

Question 5

(7 marks)

Trapezium $OABC$ is such that $\overrightarrow{AB} = 3\overrightarrow{OC}$.

The midpoints of sides OA, AB, BC and OC are P, Q, R and S .

Let $\overrightarrow{OA} = \mathbf{a}$ and $\overrightarrow{OC} = \mathbf{c}$. Use a vector method to prove that $PQRS$ is a parallelogram.

Solution
Then $\overrightarrow{OP} = \frac{1}{2}\mathbf{a}$, $\overrightarrow{OQ} = \mathbf{a} + \frac{3}{2}\mathbf{c}$. Hence $\overrightarrow{PQ} = \overrightarrow{OQ} - \overrightarrow{OP} = \mathbf{a} + \frac{3}{2}\mathbf{c} - \frac{1}{2}\mathbf{a} = \frac{1}{2}\mathbf{a} + \frac{3}{2}\mathbf{c}$. Note that $\overrightarrow{CB} = -\mathbf{c} + \mathbf{a} + 3\mathbf{c} = \mathbf{a} + 2\mathbf{c}$. Also $\overrightarrow{OS} = \frac{1}{2}\mathbf{c}$, $\overrightarrow{OR} = \mathbf{c} + \frac{1}{2}(\mathbf{a} + 2\mathbf{c}) = \frac{1}{2}\mathbf{a} + 2\mathbf{c}$. Hence $\overrightarrow{SR} = \overrightarrow{OR} - \overrightarrow{OS} = \frac{1}{2}\mathbf{a} + 2\mathbf{c} - \frac{1}{2}\mathbf{c} = \frac{1}{2}\mathbf{a} + \frac{3}{2}\mathbf{c}$. Hence $PQRS$ is a parallelogram as $\overrightarrow{PQ} = \overrightarrow{SR}$ (has a pair of equal length, parallel sides).
Specific behaviours
✓ diagram of trapezium, roughly to scale ✓ uses correct vector notation throughout ✓ vectors $\overrightarrow{OP}, \overrightarrow{OQ}$ ✓ vector $\overrightarrow{PQ}$ ✓ vectors $\overrightarrow{OS}, \overrightarrow{OR}$ ✓ vector $\overrightarrow{SR}$ ✓ conclusion

Question 6

(7 marks)

- (a) 4 different letters are chosen from the 8 in the word SCRAMBLE and then arranged to form a password. Determine how many different passwords are possible that

- (i) end in E.

(1 mark)

Solution
$n(- - - E) = 7 \times 6 \times 5 \times 1 = 210$
Specific behaviours
✓ correct number

- (ii) end in E or start with S.

(3 marks)

Solution
$n(S - - -) = 210$
$n(S - - E) = 1 \times 6 \times 5 \times 1 = 30$
$n = 210 + 210 - 30 = 390$
Specific behaviours
✓ number for D and T
✓ uses inclusion-exclusion principal
✓ correct number

- (b) Determine the number of two letter permutations that can be made using letters from the word RESCRAMBLE.

(3 marks)

Solution
8 letters - 6 singles and 2 doubles (R, E).
Both different: $8 \times 7 = 56$
Both the same: $2 \times 1 = 2$
Total permutations: $56 + 2 = 58$.
Specific behaviours
✓ breaks into exclusive cases
✓ correct calculation for each case
✓ correct number

Question 7

(7 marks)

Consider the vectors $\mathbf{a} = \begin{pmatrix} 6 \\ -7 \end{pmatrix}$, $\mathbf{b} = \begin{pmatrix} 4 \\ -3 \end{pmatrix}$ and $\mathbf{c} = \begin{pmatrix} 3 \\ -1 \end{pmatrix}$.

- (a) Determine a vector in the direction $4\mathbf{c} - \mathbf{a}$ that has the same magnitude as $\mathbf{b}$.

(3 marks)

Solution
$ \mathbf{b} = \sqrt{16 + 9} = 5$ $4\mathbf{c} - \mathbf{a} = \begin{pmatrix} 12 \\ -4 \end{pmatrix} - \begin{pmatrix} 6 \\ -7 \end{pmatrix} = \begin{pmatrix} 6 \\ 3 \end{pmatrix}$ $\left \begin{pmatrix} 6 \\ 3 \end{pmatrix} \right = \sqrt{36 + 9} = \sqrt{45} = 3\sqrt{5}$ Hence required vector is $\frac{5}{3\sqrt{5}} \begin{pmatrix} 6 \\ 3 \end{pmatrix} = \sqrt{5} \begin{pmatrix} 2 \\ 1 \end{pmatrix}$
Specific behaviours
✓ magnitude of $\mathbf{b}$ ✓ direction vector ✓ correct vector (simplification of surds optional)

- (b) Given that $\mathbf{a} = \lambda\mathbf{b} + \mu\mathbf{c}$, determine the value of λ and the value of μ .


(4 marks)

Solution
$\lambda \begin{pmatrix} 4 \\ -3 \end{pmatrix} + \mu \begin{pmatrix} 3 \\ -1 \end{pmatrix} = \begin{pmatrix} 6 \\ -7 \end{pmatrix}$ Equating i and j coefficients: $4\lambda + 3\mu = 6$ $-3\lambda - \mu = -7$ Hence $4\lambda + 3\mu = 6$ $-9\lambda - 3\mu = -21$ $-5\lambda = -15 \Rightarrow \lambda = 3$ $\mu = -3(3) + 7 = -2$ $\lambda = 3, \quad \mu = -2$
Specific behaviours
✓ equation using i -coefficients ✓ equation using j -coefficients ✓ value of λ ✓ value of μ

Question 8

(7 marks)

In the diagram shown, A, B and C lie on a circle.
The tangent at A and secant BC intersect at P .
Point Q lies on BC so that AQ bisects $\angle CAB$.



- (a) Show that $\angle PAQ = \angle PQA$.

(3 marks)

Solution
Given $\angle CAQ = \angle BAQ = \beta$. Let $\angle BAP = \alpha$, so that $\angle PAQ = \alpha + \beta$. $\angle ACB = \angle BAP = \alpha$ (alternate segment) $\angle PQA = \alpha + \beta$ (sum of opposite interior angles) Hence $\angle PAQ = \angle PQA$ as required.
Specific behaviours
✓ uses alternate segment theorem ✓ uses triangle properties ✓ logical explanation

- (b) Given that $BQ = 5$ cm and $CQ = 13$ cm, determine the length of AP .

(4 marks)

Solution
$AP^2 = PB \times PC$ (tangent secant theorem) Let $AP = PQ = x$ (isosceles triangle) so that $PB = x - 5$ and $PC = x + 13$. Then $x^2 = (x - 5)(x + 13)$ $x^2 = x^2 + 8x - 65$ $8x = 65$ $x = AP = \frac{65}{8} \text{ cm } (= 8.125)$
Specific behaviours
✓ uses intersecting secants theorem ✓ uses isosceles triangles to express required lengths ✓ forms equation ✓ correct length

Supplementary page

Question number: _____

© 2020 WA Exam Papers. Rossmoyne Senior High School has a non-exclusive licence to copy and communicate this document for non-commercial, educational use within the school. No other copying, communication or use is permitted without the express written permission of WA Exam Papers. SN085-151-7.